

BISACODYL - bisacodyl suppository

Geritrex Corp.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Bisacodyl Suppositories

Drug Facts

Active ingredient (in each suppository)	Purpose
Bisacodyl USP 10 mg	Stimulant laxative

Uses

relieves occasional constipation.

this product generally produces a bowel movement in 15 minutes to 1 hour.

Directions

tear off one suppository from the strip.

tear and peel the wrapper in downward motion.

continue tearing downward to almost the full length of the suppository (see illustration at the right)

gently remove the suppository from the wrapper.

lie on your side and push suppository. with pointed end first, high into the rectum so it will not slip out

retain it for 15 to 20 minutes.

Do not exceed recommended dose

if suppository seems soft, place in refrigerator for a short time before use

Adults and children 12 years and over 1 suppository once daily.

Children under 12 years Ask a doctor.

Adults and children 12 years and over 1 suppository once daily.

Children under 12 years Ask a doctor.

WARNINGS

For rectal use only.

Do not use when abdominal pain, nausea, or vomiting are present.

Ask a doctor before use if you have a sudden change in bowel habits that lasts over a period of 2 weeks.

When using this product you may experience. rectal burning. mild cramps.abdominal discomfort. faintness.

stop use and ask a doctor if you have rectal bleeding. you fail to have a bowel movement after using a laxative. These may be signs of a serious condition.

you need to use a laxative for more than one week.

if pregnant or breast-feeding, ask a health professional before use.

INACTIVE INGREDIENT

hydrogenated vegetable oil

STORAGE

store at temperatures below 30°C (86°F)

KEEP OUT OF REACH OF CHILDREN

If swallowed, get medical help or contact a poison control center right away.



Enter section text here

BISACODYL

bisacodyl suppository

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54162-006	
Route of Administration	RECTAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
BISACODYL (UNII: 10 X0709 Y6I) (BISACODYL - UNII:10 X0709 Y6I)		BISACODYL	0.01 g in 2 g	
Inactive Ingredients				
Ingredient Name			Strength	
HYDROGENATED SOYBEAN OIL (UNII: A2M9 1M9 18C)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54162-006-12	2 g in 1 BOX, UNIT-DOSE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part334	02/23/2012		

Labeler - Geritrex Corp. (112796248)

Revised: 2/2012

Geritrex Corp.